

Further Developments in Finite Fibonomial Calculus

Ewa Krot

Institute of Computer Science, Białystok University
 PL-15-887 Białystok, ul.Sosnowa 64, POLAND
 e-mail: ewakrot@wp.pl

February 1, 2008

Abstract

Primary definitions, notation and general observations of Finite Fibonomial Operator Calculus (FFOC) are presented. A.K.Kwaśniewski's combinatorial interpretation of Fibonomial coefficients by the use of Fibonacci cobweb poset is given. Some elements of Incidence Algebra of Fibonacci cobweb poset $\mathbf{P}$ are defined.

1 Introduction

Finite Fibonomial Operator Calculus (FFOC) is a special case of Extended Finite Operator Calculus (FOC). The "Calculus of Sequences" was started by Ward in 1936 [1] and considered by many authors after him. The very foundations of FOC were given by O.V.Viskov, G. Markowsky and finally by the author of [8, 2, 3]. The idea of FFOC due to A.K.Kwaśniewski [2] (see Example 2.1) is now continued [4] by the present author.

2 Primary Definitions, notation and general observations of FFOC

FFOC ows on the famous Fibonacci sequence $F = \{F_n\}_{n \geq 1}$ introduced by Leonardo Fibonacci [Pisano] in 1202 ("Liber Abaci") defined by the recurrence relation:

$$\begin{cases} F_0 = 0, F_1 = F_2 = 1 \\ F_{n+2} = F_{n+1} + F_n \quad n \geq 1 \end{cases} \quad (1)$$

or equivalently by the formula (Binet form):

$$F_n = \frac{1}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1 - \sqrt{5}}{2} \right)^n, \quad n \geq 0. \quad (2)$$

Let us define for above sequence what follows:

(1) F -factorial:

$$n_F! \equiv F_n! \equiv F_n F_{n-1} \dots F_2 F_1, \quad F_0! = 1.$$

(2)

$$n_F^k \equiv n_F (n-1)_F (n-2)_F \dots (n-k+1)_F \equiv F_n F_{n-1} F_{n-2} \dots F_{n-k+1}$$

(3) and finally F -binomial (Fibonomial) coefficients:

$$\binom{n}{k}_F = \frac{F_n!}{F_k! F_{n-k}!} = \frac{n_F^k}{k_F!}, \quad \binom{n}{0}_F = 1. \quad (3)$$

In the following we shall consider the algebra $\mathbf{P}$ of polynomials over the field $\mathbf{K}$ of characteristic zero.

The main object of FFOC is the F -derivative defined below.

Definition 2.1. *The F -derivative is the linear operator $\partial_F : \mathbf{P} \rightarrow \mathbf{P}$ such that $\partial_F x^n = F_n x^{n-1}$ for $n \geq 0$.*

Definition 2.2. *The F -translation operator is the linear operator $E^y(\partial_F) : \mathbf{P} \rightarrow \mathbf{P}$ of the form:*

$$E^y(\partial_F) = \exp_F \{y \partial_F\} = \sum_{k \geq 0} \frac{y^k \partial_F^k}{F_k!}, \quad y \in \mathbf{K}$$

and

$$\forall_{p \in \mathbf{P}} \quad p(x +_F y) = E^y(\partial_F) p(x) \quad x, y \in \mathbf{K}.$$

Definition 2.3. *A linear operator $T : \mathbf{P} \rightarrow \mathbf{P}$ is said to be ∂_F -shift invariant iff*

$$\forall_{y \in \mathbf{K}} \quad [T, E^y(\partial_F)] = T E^y(\partial_F) - E^y(\partial_F) T = 0$$

We shall denote by Σ_F the algebra of F -linear ∂_F -shift invariant operators.

Definition 2.4. Let $Q(\partial_F)$ be a formal series in powers of ∂_F and $Q(\partial_F) : \mathbf{P} \rightarrow \mathbf{P}$. $Q(\partial_F)$ is said to be ∂_F -delta operator iff

- (a) $Q(\partial_F) \in \Sigma_F$
- (b) $Q(\partial_F)(x) = \text{const} \neq 0$.

One can show that every ∂_F -delta operator reduces degree of any polynomial by one. For every ∂_F -delta operator $Q(\partial_F)$ there exists the uniquely determined ∂_F -basic polynomial sequence $\{q_n(x)\}_{n \geq 0}$, ($\deg q_n(x) = n$), satisfying conditions:

- (1) $q_0(x) = 1$;
- (2) $q_n(0) = 0$, $n \geq 1$;
- (3) $Q(\partial_F)q_n(x) = F_n q_{n-1}(x)$, $n \geq 0$.

Theorem 2.1. (First Expansion Theorem)

Let $T \in \Sigma_F$ and let $Q(\partial_F)$ be a ∂_F -delta operator with ∂_F -basic polynomial sequence $\{q_n\}_{n \geq 0}$. Then

$$T = \sum_{n \geq 0} \frac{a_n}{F_n!} Q(\partial_F)^n; \quad a_n = [Tq_n(x)]_{x=0}.$$

Theorem 2.2. (Isomorphism Theorem)

Let $\Phi_F = \mathbf{K}_F[[t]]$ be the algebra of formal $\exp_F$ series in $t \in \mathbf{K}$, i.e.:

$$f_F(t) \in \Phi_F \quad \text{iff} \quad f_F(t) = \sum_{k \geq 0} \frac{a_k t^k}{F_k!} \quad \text{for} \quad a_k \in \mathbf{K},$$

and let the $Q(\partial_F)$ be a ∂_F -delta operator. Then $\Sigma_F \approx \Phi_F$. The isomorphism $\phi : \Phi_F \rightarrow \Sigma_F$ is given by the natural correspondence:

$$f_F(t) = \sum_{k \geq 0} \frac{a_k t^k}{F_k!} \xrightarrow{\text{into}} T_{\partial_F} = \sum_{k \geq 0} \frac{a_k}{F_k!} Q(\partial_F)^k.$$

Remark 2.1. In the algebra Φ_F the product is given by the fibonomial convolution, i.e.:

$$\left(\sum_{k \geq 0} \frac{a_k}{F_k!} x^k \right) \left(\sum_{k \geq 0} \frac{b_k}{F_k!} x^k \right) = \left(\sum_{k \geq 0} \frac{c_k}{F_k!} x^k \right)$$

where

$$c_k = \sum_{l \geq 0} \binom{k}{l}_F a_l b_{k-l}.$$

Definition 2.5. A polynomial sequence $\{s_n\}_{n \geq 0}$ is called the sequence of Sheffer F -polynomials of the ∂_F -delta operator $Q(\partial_F)$ iff

- (1) $s_0(x) = \text{const} \neq 0$;
- (2) $Q(\partial_F)s_n(x) = F_n s_{n-1}(x)$; $n \geq 0$.

It can be shown that $\{s_n\}_{n \geq 0}$ is the sequence of Sheffer F -polynomials of ∂_F -delta operator $Q(\partial_F)$ with ∂_F -basic polynomial sequence $\{q_n\}_{n \geq 0}$ iff there exists an invertible $S \in \Sigma_F$ such that $s_n(x) = S^{-1}q_n(x)$ for $n \geq 0$. We shall refer to a given labeled by ∂_F -shift invariant invertible operator S Sheffer F -polynomial sequence $\{s_n\}_{n \geq 0}$ as the sequence of Sheffer F -polynomials of the ∂_F -delta operator $Q(\partial_F)$ relative to S .

3 Some examples of F -polynomials

The Sheffer F -polynomials form a wide class of new polynomial sequences. Some examples of them are given below:

- (1) ∂_F -basic polynomials of the operator $\Delta_F = E^1(\partial_F) - I$:

$$q_n(x) = \frac{F_n!}{n!} \hat{x}_F(E^{-1}(\partial_F)\hat{x})^{n-1}[1]$$

where $\hat{1}_F^{-1}\hat{x}_F = \hat{x}$ for $(\hat{x}p)(x) = xp(x)$, $p \in \mathbf{P}$.

- (2) ∂_F -basic polynomials of the operator $\nabla_F = I - E^{-1}(\partial_F)$:

$$q_n(x) = \frac{F_n!}{n!} \hat{x}_F(E(\partial_F)\hat{x})^{n-1}[1].$$

- (3) ∂_F -basic polynomials of F -Abel operator $A(\partial_F) = \partial_F E^a(\partial_F) = \sum_{k \geq 0} \frac{a^k}{F_k!} \partial_F^{k+1}$:

$$A_{n,F}^{(a)}(x) = \frac{F_n}{n} \sum_{k=0}^{n-1} \binom{n-1}{k}_F (-an)^k \frac{n-k}{F_{n-k}} x^{n-k}.$$

Here are the first examples of Abel F -polynomials of order a :

$$A_{0,F}^{(a)}(x) = 1$$

$$A_{1,F}^{(a)}(x) = x$$

$$A_{2,F}^{(a)}(x) = x^2 + ax$$

$$A_{3,F}^{(a)}(x) = x^3 - 4ax^2 + 6a^2x$$

$$A_{4,F}^{(a)}(x) = x^4 - 9ax^3 + 48a^2x^2 + 48a^3x$$

$$A_{5,F}^{(a)}(x) = x^5 - 20ax^4 + 225a^2x^3 - 750a^3x^2 + 625a^4x$$

$$A_{6,F}^{(a)}(x) = x^6 - 40ax^5 + 960a^2x^4 - 6480a^3x^3 + 17280a^4x^2 - 10368a^5x$$

$$A_{7,F}^{(a)}(x) = x^7 - 78ax^6 + 3640a^2x^5 - 50960a^3x^4 + 267540a^4x^3 - 499408a^5x^2 + 218491a^6x$$

$$A_{8,F}^{(a)}(x) = x^8 - 147ax^7 + 131004a^2x^6 - 349440a^3x^5 + 3727360a^4x^4 - 13418496a^5x^3 + 17891328a^6x^2 - 5505024a^7x.$$

(4) ∂_F -basic polynomials of F -Laguerre operator $L(\partial_F) = \frac{\partial_F}{\partial_F - I} = \sum_{k \geq 0} \partial_F^{k+1}$:

$$L_{n,F}(x) = \frac{F_n}{n} \sum_{k=1}^n (-1)^k \frac{F_n!}{F_k!} \binom{n-1}{k-1}_F \frac{k}{F_k} x^k.$$

Here are the few first of them:

$$L_{0,F}(x) = 1$$

$$L_{1,F}(x) = -x$$

$$L_{2,F}(x) = x^2 - \frac{1}{2}x$$

$$L_{3,F}(x) = -x^3 + \frac{8}{3}x^2 - \frac{4}{3}x$$

$$L_{4,F}(x) = x^4 - \frac{27}{4}x^3 + 18x^2 - \frac{9}{2}x$$

$$L_{5,F}(x) = -x^5 + 20x^4 - 135x^3 + 180x^2 - 30x$$

$$L_{6,F}(x) = x^6 - \frac{160}{3}x^5 + \frac{3200}{3}x^4 - 3600x^3 + 3200x^2 - 320x$$

$$L_{7,F}(x) = -x^7 + \frac{13182}{7}x^6 - \frac{54080}{7}x^5 + \frac{540800}{7}x^4 - \frac{1216800}{7}x^3 + \frac{648960}{7}x^2 - \frac{40560}{7}x$$

$$L_{8,F}(x) = x^8 - \frac{3087}{8}x^7 + \frac{223587}{4}x^6 - 1498580x^5 + 9937200x^4 - 13415220x^3 + 4471740x^2 - 171990x.$$

- (5) Hermite F -polynomials are Sheffer F -polynomials of the ∂_F -delta operator ∂_F relative to invertible $S \in \Sigma_F$ of the form $S = \exp_F\{\frac{a\partial_F^2}{2}\}$:

$$H_{n,F}(x) = S^{-1}x^n = \sum_{k \geq 0} \frac{(-a)^k}{2^k F_k!} n_{F_k}^{2k} x^{n-2k}.$$

- (6) Let $S = (1 - \partial_F)^{-\alpha-1}$. The Sheffer F -polynomials of ∂_F -delta operator $L(\partial_F) = \frac{\partial_F}{\partial_F - 1}$ relative to S are Laguerre F -polynomials of order α are given by formula:

$$L_{n,F}^{(\alpha)}(x) = \sum_{k \geq 0} \frac{F_n!}{F_k!} \binom{\alpha + n}{n - k}_F (-x)^k$$

for $\alpha \neq -1$. Here are the few first of Laguerre F -polynomials of order $\alpha = 1$:

$$L_{0,F}^{(1)}(x) = 1$$

$$L_{1,F}^{(1)}(x) = -x + 1$$

$$L_{2,F}^{(1)}(x) = x^2 - 2x + 2$$

$$L_{3,F}^{(1)}(x) = -2x^3 + 6x^2 - 12x + 6$$

$$L_{4,F}^{(1)}(x) = 6x^4 - 30x^3 + 90x^2 - 90x + 30$$

$$L_{5,F}^{(1)}(x) = -30x^5 + 240x^4 - 1200x^3 + 1800x^2 - 1200x + 240$$

$$L_{6,F}^{(1)}(x) = 240x^6 - 3120x^5 + 24960x^4 - 62400x^3 + 62400x^2 - 24960x + 3120.$$

(7) Bernoullie's F -polynomials of order 1 are Sheffer F -polynomials of ∂_F -delta operator ∂_F related to invertible $S = \left(\frac{\exp_F\{\partial_F\} - I}{\partial_F} \right)^{-1}$. They are given by formula:

$$B_{n,F}(x) = \sum_{k \geq 0} \frac{1}{F_{k+1}} \binom{n}{k}_F x^{n-k}.$$

Here we give few first of them:

$$B_{0,F}(x) = 1$$

$$B_{1,F}(x) = x + 1$$

$$B_{2,F}(x) = x^2 + x + \frac{1}{2}$$

$$B_{3,F}(x) = x^3 + 2x^2 + x + \frac{1}{3}$$

$$B_{4,F}(x) = x^4 + 3x^3 + 3x^2 + x + \frac{1}{5}$$

$$B_{5,F}(x) = x^5 + 5x^4 + \frac{15}{2}x^3 + 5x^2 + x + \frac{1}{8}$$

$$B_{6,F}(x) = x^6 + 8x^5 + 20x^4 + 20x^3 + 8x^2 + x + \frac{1}{13}$$

$$B_{7,F}(x) = x^7 + 13x^6 + 52x^5 + \frac{260}{3}x^4 + 52x^3 + 13x^2 + x + \frac{1}{21}$$

$$B_{8,F}(x) = x^8 + 21x^7 + \frac{273}{2}x^6 + 364x^5 + 364x^4 + \frac{273}{2}x^3 + 21x^2 + x + \frac{1}{36}$$

$$B_{9,F}(x) = x^9 + 34x^8 + 357x^7 + 1547x^6 + \frac{12376}{5}x^5 + 1547x^4 + 357x^3 + 34x^2 + x + \frac{1}{55}.$$

Remark 3.1. It is known (from the general case, see [8, 7, 4]) that every sequence of Sheffer F -polynomials is orthogonal in respect to special inner product associated with this sequence. The problem of classical orthogonality of these polynomials is under investigation.

4 Combinatorial Content of FFOC

Let us quote some standard interpretations:

- binomial coefficients $\binom{n}{k}$ denote the number of k -element subsets of set with n elements;
- Stirling numbers of the first kind $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ denote the number of permutations of n elements containing exactly k cycles;
- Stirling numbers of the second kind $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ denote the number of partitions of n elements into k blocks;
- q-Gaussian coefficients $\binom{n}{k}_q$ denote the number of k dimensional subspaces in n -th dimensional space over Galois field $GF(q)$.

The analogues of above numbers are fibonomial coefficients defined in Section 2:

$$\binom{n}{k}_F = \frac{F_n!}{F_k! F_{n-k}!} = \frac{n_F^k}{k_F!}, \quad \binom{n}{0}_F = 1. \quad (4)$$

One can show (using some properties of Fibonacci numbers) that they are of integer type. Our question is: **Does the combinatorial (classic like) interpretation of them exist?** It was unknown until now. The combinatorial interpretation of fibonomial coefficients has been given by A.K.Kwaśniewski in [5, 6] by the use of special partially ordered set (Fibonacci cobweb poset). His idea was to start with a famous rabbits grown Fibonacci tree. Let us define a partially ordered set $\mathbf{P}$ via its Hasse diagram. It looks like rabbits tree, i.e. it consists of levels labeled by Fibonacci numbers (the n -th level consist of F_n elements). Every element of n -th level ($n \geq 1, n \in \mathbf{N}$) is in partial order relation with every element of the $(n+1)$ -th level but it's not with any element from the level in which he lies (n -th level) except from himself. We will call this poset $\mathbf{P}$ the Fibonacci cobweb poset.

We can also define it using the characteristic function ζ of the partial order relation $\leq$ in $\mathbf{P}$. It is defined for any poset as follows:

$$\zeta(x, y) = \begin{cases} 1 & \text{for } x \leq y \\ 0 & \text{otherwise} \end{cases}$$

The ζ function of $\mathbf{P}$ may be expressed by the formula:

$$\zeta(x, y) = \begin{cases} 0 & \text{if } x > y \\ 0 & (\sum_{i=1}^{k-1} F_i) + 1 \leq x, y \leq \sum_{i=1}^k F_k; \quad x \neq y; \quad k \geq 2 \\ 1 & x = y \\ 1 & \text{otherwise} \end{cases}$$

In the above the condition $(\sum_{i=1}^{k-1} F_i) + 1 \leq x, y \leq \sum_{i=1}^k F_k; \quad x \neq y; \quad k \geq 2$ means that x, y are the elements of the same (k -th) level. We can use the following property of Fibonacci numbers:

$$\sum_{i=1}^k F_k = F_{k+2} - 1; \quad k \geq 1 \quad (5)$$

to get better looking formula for ζ function:

$$\zeta(x, y) = \begin{cases} 0 & \text{if } x > y \\ 0 & F_{k+1} \leq x, y \leq F_{k+2} - 1; \quad x \neq y; \quad k \geq 3 \\ 1 & x = y \\ 1 & \text{otherwise} \end{cases} \quad (6)$$

The incidence ζ function infinite matrix representing uniquely cobweb poset $\mathbf{P}$ was given at first by A.K.Kwaśniewski in [5]. It is of the form

$$\begin{bmatrix} 1 & \dots \\ & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & & & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & & & & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & & & & & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ & & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & 1 & \dots \\ & & & & & & & & & & & & & & 1 & 0 & 0 & 0 & 1 & \dots \\ & & & & & & & & & & & & & & & 1 & 0 & 0 & 1 & \dots \\ z & e & r & o & s & & & & & & & & & & & & & & & & \dots \\ & \dots \end{bmatrix}$$

From the definition of $\mathbf{P}$ follows:

Observation 4.1. [5, 6]

The number of maximal chains starting from the root (the first level), to reach n -th level (labeled by F_n) is equal to $n_F!$.

Observation 4.2. [5, 6]

The number of maximal chains starting from the any fixed point from the level labeled by F_k to reach any point at the n -th level is equal to $n_F^{n-k} = n_F^m$, ($n = k + m$).

Let us consider the following question, (quotation from [5, 6]):

"Let us denote by P_m a subposet of P consisting of points up to m -th level points: $\bigcup_{s=1}^n \Phi_s$; Φ_s is the set of elements of the s -th level. Consider now the following behaviour of a "sub-cob" moving from any given point of the F_k level of the poset up. It behaves as it has been born right there and can reach at first F_2 points up, then F_3 points up, F_4 and so on - thus climbing up to the level $F_{k+m} = F_n$ of the poset $\mathbf{P}$. It can see- its Great Ancestor at the root F_1 -th level and potentially follow one of its own accessible finite subposet P_m . One of many P_m 's rooted at the k -th level may be found. How many?"

Observation 4.3. [5, 6]

Let $n = k + m$. The number of subposets P_m rooted at any fixed point at the level labeled by F_k and ending at the n -th level (labeled by F_n) is equal to:

$$\binom{n}{m}_F = \binom{n}{m}_f = \frac{n_F^k}{k_F!}.$$

5 The Incidence Algebra $\mathbf{I}(\mathbf{P})$

One can define the incidence algebra of $\mathbf{P}$ (locally finite partially ordered set) as follows (see [9, 10]):

$$\mathbf{I}(\mathbf{P}) = \{f : \mathbf{P} \times \mathbf{P} \longrightarrow \mathbf{R}; \quad f(x, y) = 0 \text{ unless } x \leq y\}.$$

The sum of two such functions f and g and multiplication by scalar are defined as usual. The product $H = f * g$ is defined as follows:

$$h(x, y) = (f * g)(x, y) = \sum_{z \in \mathbf{P}: x \leq z \leq y} f(x, z) * g(z, y).$$

It is immediately verified that this is an associative algebra over the real field (associative ring).

The incidence algebra has an identity element $\delta(x, y)$, the Kronecker delta. The zeta function of $\mathbf{P}$ is an element of $\mathbf{I}(\mathbf{P})$. It was expressed by δ in [6] from where we learn that:

$$\zeta = \zeta_1 - \zeta_0$$

where for $x, y \in \mathbf{N}$:

$$\zeta_1(x, y) = \sum_{k=0}^{\infty} \delta(x + k, y) \quad (7)$$

$$\zeta_0(x, y) = \sum_{k \geq 0} \sum_{s \geq 0} \delta(x, F_{s+1} + k) \sum_{r=1}^{F_s - k - 1} \delta(k + F_{s+1} + r, y). \quad (8)$$

The knowledge of ζ enables us to construct other typical elements of incidence algebra perfectly suitable for calculating number of chains, of maximal chains etc. in finite sub-posets of $\mathbf{P}$. The one of them is Möbius function indispensable in numerous inversion type formulas of countless applications. It is known that the ζ function of a locally finite partially ordered set is invertible in incidence algebra and its inversion is so called the Möbius function μ i.e.:

$$\zeta * \mu = \mu * \zeta = \delta.$$

The Möbius function μ of Fibonacci cobweb poset $\mathbf{P}$ is presented here for the first time and it's due to the present author. It can be recovered using the following recurrence formula (see [9]):

$$\begin{cases} \mu(x, x) = 1 & \text{for all } x \in \mathbf{P} \\ \mu(x, y) = - \sum_{x \leq z < y} \mu(x, z) \end{cases} \quad (9)$$

Then we get:

$$\mu(x, y) = \begin{cases} 0 & x > y \\ 1 & x = y \\ 0 & F_{k+1} \leq x, y \leq F_{k+2} - 1; x \neq y; k \geq 3 \\ -1 & F_{k+1} \leq x \leq F_{k+2} - 1 < F_{k+2} \leq y \leq F_{k+3} - 1 \\ -\prod_{l=k+1}^{n-1} (1 - F_l) & F_{k+1} \leq x \leq F_{k+2} - 1, \\ & F_{n+1} \leq y \leq F_{n+2} - 1; n > k + 1, \end{cases} \quad (10)$$

where:

- the condition $F_{k+1} \leq x, y \leq F_{k+2} - 1$; $x \neq y$; $k \geq 3$ means that x, y are different elements of k -th level;
- the condition $F_{k+1} \leq x \leq F_{k+2} - 1 < F_{k+2} \leq y \leq F_{k+3} - 1$ means that x is an element of k -th level and y is an element of $(k + 1)$ -th level;
- the condition $F_{k+1} \leq x \leq F_{k+2} - 1$, $F_{n+1} \leq y \leq F_{n+2} - 1$; $n > k + 1$ means that x is an element of k -th level and y is an element of n -th level.

So the μ function matrix has the following structure:

$$\begin{bmatrix} 1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ & 1 & -1 & -1 & 1 & 1 & 1 & -2 & -2 & -2 & -2 & -2 & 8 & 8 & 8 & 8 & 8 & \dots \\ & & 1 & 0 & -1 & -1 & -1 & 2 & 2 & 2 & 2 & 2 & -8 & -8 & -8 & -8 & -8 & \dots \\ & & & 1 & -1 & -1 & -1 & 2 & 2 & 2 & 2 & 2 & -8 & -8 & -8 & -8 & -8 & \dots \\ & & & & 1 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & 4 & 4 & 4 & 4 & 4 & \dots \\ & & & & & 1 & 0 & -1 & -1 & -1 & -1 & -1 & 4 & 4 & 4 & 4 & 4 & \dots \\ & & & & & & 1 & -1 & -1 & -1 & -1 & -1 & 4 & 4 & 4 & 4 & 4 & \dots \\ & & & & & & & 1 & 0 & 0 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & \dots \\ & & & & & & & & 1 & 0 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & \dots \\ & & & & & & & & & 1 & 0 & 0 & -1 & -1 & -1 & -1 & -1 & \dots \\ & & & & & & & & & & 1 & 0 & -1 & -1 & -1 & -1 & -1 & \dots \\ & & & & & & & & & & & 1 & -1 & -1 & -1 & -1 & -1 & \dots \\ & & & & & & & & & & & & 1 & 0 & 0 & 0 & 0 & \dots \\ & & & & & & & & & & & & & 1 & 0 & 0 & 0 & \dots \\ & & & & & & & & & & & & & & 1 & 0 & 0 & \dots \\ & & & & & & & & & & & & & & & 1 & 0 & \dots \\ & & & & & & & & & & & & & & & & 1 & \dots \\ & & & & & & & & & & & & & & & & & \dots \\ & & & & & & & & & & & & & & & & & \dots \\ z & e & r & o & s & & & & & & & & & & & & & \dots \\ & & & & & & & & & & & & & & & & & \dots \end{bmatrix}$$

Acknowledgements

I would like to thank to Prof. A.K.Kwaśniewski for his remarks and guideness.

References

- [1] M.Ward: *A calculus of Sequences*, Amer. J. Math.**58**(1936)255-266.

- [2] A.K.Kwaśniewski: *On Simple Characterizations of Sheffer ψ -polynomials and Related Propositions of the Calculus of Sequences*, Bulletin de la Soc. des Sciences et de Lettres de Lodz, 52, Ser.Rech.Deform. **36**(2002)45-65 (see also ArXiv; Paper:math.CO/0312397)
- [3] A.K.Kwaśniewski: *Main Theorems of Extended Finite Operator Calculus*, Integral Transforms and Special Functions Vol 14, No 6,(2003)499-516
- [4] E.Krot: *An Introduction to Finite Fibonomial Calculus*, Inst.Comp.Sci.UwB/Preprint#50/October/2003 submitted to publication
- [5] .K.Kwaśniewski:*Information on combinatorial Interpretation of FibonomialCoefficients*, Bull.Soc.Lett.Lodz.Ser.Rech.Deform. 42(2003) p.39-41 ArXiv:math.CO/0402291 v1 22 Feb 2004
- [6] A.K.Kwaśniewski:*More on Combinatorial Interpretation of Fibonomial Coefficients*,ArXiv:math.CO/0402344 v1 18 Feb 2004
- [7] E.Krot: *ψ -extensions of q -Hermite and q -Laguerre Polynomials - properties and principal statements*, Czech. J. Phys. Vol.**51**(2001)No12, p.1362-1367.
- [8] A. K. Kwaśniewski: *Towards ψ -Extension of Rota's Finite Operator Calculus*, Rep. Math. Phys. **47**, 305 (2001)p.305-342 ArXiv: math.CO/0402078 2004
- [9] G.C.Rota: *On the Foundations of Combinatorial Theory I: Theory of MöFunctions*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, Vol.2 (1964)p. 340-368
- [10] E.Spiegel, J.O'Donnell: Incidence Algebras, Marcel Dekker, Inc. Basel 1997

AMS Classification numbers: 11C08, 11B37, 47B47